

Michigan Quality Improvement Consortium Guideline General Principles for the Diagnosis and Management of Asthma

The following guideline recommends general principles and key clinical activities for the diagnosis and management of asthma.

Eligible Population	Key Components	Recommendation and Level of Evidence
<p>Children and adults with the following:</p> <ul style="list-style-type: none"> • Wheezing • History of cough (worse particularly at night), recurrent wheeze, recurrent difficulty in breathing, recurrent chest tightness • Symptoms occur or worsen in the presence of exercise, viral infection, inhalant allergens, irritants, changes in weather, strong emotional expression (laughing or crying hard), stress, menstrual cycles • Symptoms occur or worsen at night, awakening the patient 	Diagnosis and management goals	<ul style="list-style-type: none"> • Detailed medical history and physical exam to determine that symptoms of recurrent episodes of airflow obstruction are present • Use spirometry in all patients > 5 years of age to determine that airway obstruction is at least partially reversible [C]. • Consider alternative causes of airway obstruction. <p>Goals of therapy are to achieve control by [A]:</p> <ul style="list-style-type: none"> • Reducing impairment (prevent chronic symptoms, minimize need for rescue therapy with short-acting beta₂-agonists (SABA), maintain near-normal lung function and activity levels) • Reducing risk (prevent exacerbations, minimize need for emergency care or hospitalization, prevent loss of lung function or prevent reduced lung growth in children, have minimal or no adverse effects of therapy)
	Assessment and monitoring	<ul style="list-style-type: none"> • Assess asthma severity to initiate therapy. (Use severity classification chart, assessing both domains of impairment [B] and risk [C] to determine initial treatment.) • Assess asthma control to monitor and adjust therapy [B]. (Use asthma control chart, assessing both domains of impairment and risk to determine if therapy should be maintained or adjusted. (Step up if necessary; step down if possible.)) • Obtain lung function measures by spirometry at least every 1-2 years [B], more frequently for not well-controlled asthma. • Schedule follow-up care: In general, consider scheduling patients at 2 – to 6-week intervals while gaining control [D]; at 1- to 6-month intervals, depending on step of care required or duration of control, to monitor if sufficient control is maintained; at 3-month intervals if a step-down in therapy is anticipated [D]. • Assess asthma control, medication technique, written asthma action plan, patient adherence and concerns at every visit.
	Education	<ul style="list-style-type: none"> • Provide self-management education [A]. Teach and reinforce: self-monitoring to assess control and signs of worsening asthma (either symptom or peak flow monitoring) [B]; using written asthma action plan (review differences between long-term control and quick-relief medication); taking medication correctly (inhaler technique and use of devices); avoiding environmental and occupational factors that worsen asthma. • Tailor education to literacy level of patient; integrate education into all points of care; appreciate potential role of patient’s cultural beliefs and practices in asthma management [C]. • Develop written action plan in partnership with patient [B].
	Control environmental factors and comorbid conditions	<ul style="list-style-type: none"> • Recommend measures to control exposures to allergens and pollutants or irritants that make asthma worse [A]. • Consider allergen immunotherapy for patients with persistent asthma and when there is clear evidence of a relationship between symptoms and exposure to an allergen to which the patient is sensitive [B]. • Treat comorbid conditions (e.g., allergic bronchopulmonary aspergillosis [A], gastroesophageal reflux [B], obesity [B], obstructive sleep apnea [D], rhinitis and sinusitis [B], chronic stress or depression) [D]. • Inactivated influenza vaccine for all patients over 6 months of age [A] unless contraindicated
	Medications	<ul style="list-style-type: none"> • Select medication and delivery devices to meet patient’s needs. • Use a stepwise approach to pharmacologic therapy to gain and maintain asthma control [A]. (See age-specific guidelines.) • Inhaled corticosteroids (ICS) are the most effective long-term control therapy [A]. Optimize ICS use before advancing to other therapies. When choosing among treatment options, consider patient’s impairment and risk, history of response to medication, willingness and ability to use medication.
	Referral	<ul style="list-style-type: none"> • Refer to an asthma specialist for consultation or comanagement if there are difficulties achieving or maintaining control (See age-specific guidelines.); immunotherapy or omalizumab is considered; additional testing is indicated; or if the patient required 2 bursts of oral systemic corticosteroids in the past year or a hospitalization [D].

Levels of Evidence for the most significant recommendations: A=randomized controlled trials; B=controlled trials, no randomization; C=observational studies; D=opinion of expert panel

This guideline lists core management steps. It is based on the 2007 National Asthma Education and Prevention Program Expert Panel Report 3, Guidelines for the Diagnosis and Management of Asthma, National Heart, Lung and Blood Institute (www.nhlbi.nih.gov). Individual patient considerations and advances in medical science may supersede or modify these recommendations.